

Bicuspid Aortic Valve Characteristics in Children

Habibollah Yadollahi Farsani, MD and Akbar Shahmohammadi Beni*, MD

Abstract

Background- Bicuspid aortic valve (BAV) is the most common congenital heart disease and the most common malformation of aortic valve. In BAV, there are two cusps instead of three cusps in the aortic valve. The objectives of this study were the determination of the aortic root dilatation and other anatomic and hemodynamic characteristics and abnormalities of BAV.

Methods- Thirty patients and 30 control subjects were evaluated. Aortic root dimensions were measured via two-dimensional echocardiography (2-D echo) at 4 levels, including the aortic valve annulus, sinuses of Valsalva, sinotubular junction (STJ) and proximal ascending aorta (AAO). Hemodynamic data and anatomic characteristics were measured using 2D and Doppler-echo. All the findings were matched and indexed for body surface area (BSA) and were compared with the matched data of the control subjects. Clinical and demographic findings of BAV were also determined and collected through a questionnaire.

Results- Among the patients, 70% were male and the mean age and weight of the patients were 7.5 years and 22.13 kg, respectively. 86.66% of the patients had systolic ejection murmur (SEM), 76.66% systolic ejection click (SEC) and 10% had chest pain. Other congenital heart diseases (CHD) were found in 26.96% of the patients, including coarctation of the aorta (CoA) in 23% of the cases. Matched mean anatomic aortic valve area (AAVA) was $2.05\text{cm}^2/\text{m}^2$, and matched mean effective aortic valve area (EAVA) was $1.41\text{cm}^2/\text{m}^2$ BSA. Maximum aortic valve pressure gradient (PG max) in systole was 56.56mmHg. Forty percent of the patients had aortic stenosis (AS): mild AS in 16.66%, moderate AS in 13.33% and intermediate AS in 10%. Prevalence of aortic insufficiency (AI) was 36.68%. When the data were compared with the control subjects, all the patients showed a meaningful larger aortic root dimension at all 4 levels (P values are presented in Table IV). Aortic root dilation was at the level of the annulus, sinuses of Valsalva, STJ and proximal AAO in 6.25%, 4.75%, 10.20% and 10.13%, respectively.

Conclusion- These findings support the hypothesis that BAV and aortic root dilation may reflect a common developmental defect. AS and AI are common in BAV. Similar to other obstructive defects of the left heart, BAV is significantly more common in males. Because murmurs and clicks are common in BAV even without AS or AI, all patients with a heart murmur and/or click must be evaluated for BAV (*Iranian Heart Journal 2008; 9 (1):40 -46*).

Key words: bicuspid aortic valve ■ congenital heart disease ■ aortic root dilation ■ children

The bicuspid aortic valve (BAV) is the most common congenital heart disease (CHD), and is found in 1-2% of the general population. BAV is the most common malformation of the aortic valve.

Patients with BAV are susceptible to infective endocarditis (moderate risk). As the most common CHD and due to its serious complications, BAV must be diagnosed as soon as possible.

Received Apr. 21 2006; Accepted for publication Apr 24, 2008.

From the Department of Pediatrics, Hajar Hospital, Parastar (Varzesh) Street, Shahrekord, and the *Department of Pediatric Cardiology, Shaheed Rajaie Cardiovascular Medical Center, Tehran, Iran.

Corresponding author: Habibollah Yadollahi Farsani, MD, Pediatric Cardiologist, Hajar Hospital, Parastar (Varzesh) Street, Post code: 88167-54633, Shahrekord, Iran. Tel (Hospital): 0381-2223350, 2220016 Fax: 0381-2243715 E-mail: Hooman_4028 @ Yahoo.com

In this study, we describe our findings in patients referred to the pediatric clinic for medical advice.

In BAV, there are 2 cusps instead of 3 cusps in the aortic valve (Fig. 1).




Fig.1. Echocardiographic examinations in parasternal short axis view. Left panel; diastole; the presence of a raphe at 1 o'clock simulates a normal tricuspid aortic valve.

Right panel: midsystole; it is clearly demonstrated that there are only two cusps.

Methods

This was a case-series study with a sample volume of 30 cases, and the duration of the study was 12 months. All the patients were referred to the outpatient pediatric cardiology clinic of Shaheed Rajaie Cardiovascular Medical Center during 2003-2004 (12 months). They were visited by pediatric cardiologists and their assistants. All the demographic data were obtained from the patients and their parents. The patients were examined, and then all the collected data and clinical findings were recorded in special questionnaires after the documentation of BAV via echocardiography. For the estimation of AAVA and EAVA, we used a continuity equation:

$EAVA = LVOT_{CSA} \times LVOT_{VTI} / AV_{VTI}$ and $\pi d^2/4$, where EAVA= effective aortic valve area, LVOT= left ventricle outflow tract, AV= aortic valve, CSA= cross-sectional area, VTI= velocity time integral (estimated by Doppler echo), d= diameter of aortic valve annulus and LVOT and $\pi=3.14$. Area unit was cm^2 , and VTI unit was cm/s (second). For data analysis, we also used T-test and χ^2 . This study has been done for the first time in Iran.




Fig. 2. Diagrammatic representation of the aortic root with the sites of measurement: 1, aortic annulus; 2, sinuses of Valsalva; 3, supra-aortic ridge; 4, proximal ascending aorta. Ao, aorta; LA, left atrium; LV, left ventricle.

Results

70% of the patients were male (M), and 30% female (F). The mean age and weight (wt) of the patients were 7.5 years and 22.13 kg, respectively. About 87% of the patients had systolic ejection murmur (SEM), and 76.66% had systolic ejection click (SEC) and about 10% had chest pain (CP). Other congenital heart diseases (CHD) were found in 26.96% of the patients, including coarctation of the aorta (CoA) 23%, ventricular septal defect (VSD) 6.66%, atrial septal defect (ASD) 3.33% and congenital complete heart block (CHB) 3.33% who underwent permanent pacemaker (PPM) implantation (Table I, Figs. 3, 4).

TableI. Demographic variables arranged according to age and number.

Patient ^a	No. ^b	Age			Sex		Wt (kg) ^k	HT (Cm) ^l	BSA ^m (m ²)	MM ^q		CP ^z	Click ^y	Other CHD	Kind of CHD ^w
		D ^c	M ^d	Y ^e	F ^f	M ^g				SM ^p	DM ^s				
K-B	1	17	-	-	-	+	3	50	0.20	+	-	-	-	+	ASD, CoA
M-Z	2	-	10	-	-	+	8	72	0.45	+	-	-	-	+	VSD, CoA
K-F	3	-	11	-	+	-	8	71	0.44	+	-	-	-	-	-
Z-S	4	-	6	1	+	-	10	88	0.50	+	-	-	+	-	-
A-S	5	-	-	3	-	+	15	110	0.66	+	+	-	+	-	-
M-M	6	-	4	3	-	+	12	92	0.55	+	-	-	+	-	-
B-B	7	-	6	4	+	-	19	124	0.82	-	+	-	+	-	-
H-B	8	-	-	5	-	+	20	122	0.82	+	-	-	+	-	-
F-Gh	9	-	4	5	+	-	18	116	0.76	+	+	-	+	-	-
M-K	10	-	-	5	+	-	15	112	0.66	-	-	-	-	+	CoA
M-M	11	-	9	6	-	+	17	110	0.70	+	-	-	+	-	-
M-T	12	-	-	7	-	+	17	115	0.73	+	+	-	+	-	-
S-M	13	-	3	7	+	-	20	126	0.82	+	+	-	+	-	-
M-K	14	-	5	7	-	+	23	140	0.92	+	-	-	-	-	-
S-S	15	-	2	7	+	-	16	115	0.70	+	+	-	+	-	-
A-A	16	-	10	7	-	+	22	129	0.86	+	-	-	+	-	-
H-A	17	-	10	8	-	+	25	142	0.96	+	-	-	+	-	-
M-N	18	-	10	8	-	+	23	124	0.88	+	-	-	+	+	CoA
H-H	19	-	11	8	-	+	23	130	0.90	+	+	-	+	-	-
H-M	20	-	11	8	-	+	20	120	0.82	+	+	-	+	-	-
A-KH	21	-	-	9	-	+	18	120	0.80	+	-	-	+	-	-
R-S	22	-	-	10	-	+	23	126	0.88	+	-	-	+	-	-
R-A	23	-	2	10	-	+	30	154	1.10	-	-	-	-	-	-
M-V	24	-	-	11	-	+	35	161	1.22	-	-	-	-	+	CHB+PPM
N-B	25	-	2	12	+	-	35	170	1.25	+	+	+	+	-	-
A-S	26	-	-	13	+	-	30	160	1.10	+	-	-	-	+	CoA
A-A	27	-	-	13	-	+	37	150	1.24	+	-	+	+	-	-
H-A	28	-	6	13	-	+	35	165	1.20	+	+	-	+	-	-
S-A	29	-	-	14	-	+	55	170	1.60	+	-	+	+	+	CoA
M-S	30	-	2	14	-	+	32	150	1.12	+	+	-	+	+	CoA-VSD
Consi-deration	30	17	124	218	9	21	22.13 (3-55)	124.46 (50-170)	0.85 (0.20-1.60)	26	11	3	23	8	CoA=7 VSD=2 ASD=1 CHB=1
	Total	7.5 y (17d-14y) Mean			30.00%	70.00%	Mean			86.66%	36.66%	10.00%	76.66%	26.66%	

a: first and last initial; b: number of patient’s record; c: day ; d: month; e: year; f: female; g: male; k: weight in kilograms; l: height in centimeters; m: body surface area in square meters; q: murmur; p: systolic murmur; s: diastolic murmur; z: chest pain; ASD: atrial septal defect; CHB: complete heart block; CoA: coarctation of aorta; PPM: permanent pacemaker; VSD: ventricular septal defect




Fig. 3. Prevalence of clinical findings




Fig. 5. EAVA/AAVA ratio.




Fig. 4. Prevalence of associated CHD

Matched mean anatomic aortic valve area (AAVA) was $2.05 \text{ cm}^2/\text{m}^2$, and matched mean effective aortic valve area (EAVA) was $1.41 \text{ cm}^2/\text{m}^2$ BSA (body surface area). EAVA: AAVA ratio was about 0.68, suggesting incomplete aortic valve opening during systole (Fig. 5).

Maximum pressure gradient (PG_{max}) across the aortic valve was 56.56 mmHg in systole. Among the patients, 40% had aortic stenosis (AS), of which 16.66% was mild, 13.33% was moderate and 10% was intermediate AS. There was no case with severe AS. Prevalence of aortic insufficiency (AI) was 36.68%, all of which were mild (1^+) to moderate (2^+). When the collected echocardiographic data (Tables II, III) were compared with the control subjects (Table IV), all the patients showed meaningfully larger aortic root dimensions at the aortic valve annulus, sinuses of Valsalva, STJ and proximal AAO (Fig. 2, see Table IV P-values). The percents of the enlargement of the 4 above-mentioned diameters were 6.35% for the annulus, 4.75% for sinuses of Valsalva, 10.20% for sinotubular junction (STJ) and 10.13% for proximal ascending aorta (AAO). All the echocardiographic data were collected and summarized in Tables I-IV.

Table II. Echocardiographic variable arranged according to age and number.

1	2	3	4	5	6	7	8	9	10	File No.	Variables
3.15	2.04	1.89	2.57	1.05	2.54	2.24	2.27	1.22	1.62		Indexed-AAVA Cm ² /m ²
3.15	1.43	0.90	0.81	1.15	2.54	2.12	1.51	1.51	1.56		Indexed-EAVA Cm ² /m ²
1.00	0.70	0.48	0.31	1.00	1.00	0.93	0.66	1.00	0.96		EAVA/AAVA Ratio
3.15	1.57	1.95	3.28	2.25	3.63	2.35	3.06	1.60	1.93		Indexed-LVOTSA Cm ² /m ²
1.56	1.50	2.59	3.19	1.99	1.22	1.42	2.11	3.43	1.69		V _{max} - M/s
1.09	0.93	1.52	2.06	1.37	0.84	1.09	1.27	2.29	1.18		V _{mean} - M/s
13.43	9.01	26.87	40.73	15.83	5.96	8.16	17.86	47.11	11.50		P _{max} - mmHg
6.31	4.52	12.50	21.25	9.26	3.34	5.52	8.73	29.39	6.56		P _{mean} - mmHg
21.10	20.42	38.75	48.38	34.27	20.22	26.67	36.45	55.55	32.84		VTI-Cm
1.74	1.52	1.05	0.68	0.93	0.83	1.40	1.06	3.23	1.48		V _{max} - M/s
1.07	0.91	0.76	0.57	0.79	0.70	1.07	0.71	2.18	1.04		V _{mean} - M/s
11.77	9.38	4.45	2.79	3.50	2.78	7.79	4.52	41.72	8.83		P _{max} - mmHg
6.20	4.49	2.60	2.26	2.92	2.29	5.55	2.57	24.59	5.26		P _{mean} - mmHg
19.92	18.58	18.04	12.01	17.57	14.25	24.11	18.09	52.33	26.47		VTI-Cm
0.93	0.93	1.05	1.46	1.38	1.59	1.57	1.79	1.25	1.28		LVOT
0.66	0.66	1.03	1.28	0.94	1.38	1.57	1.54	1.09	1.17		AV-Ann
0.74	1.51	1.21	1.71	1.85	1.72	2.48	2.23	1.64	1.55		Vals-Sin
0.66	1.00	1.02	1.22	0.99	1.22	1.93	1.74	1.19	1.18		STJ
1.09	1.60	1.33	1.85	1.35	1.38	1.60	1.98	1.92	1.60		AAO
11	12	13	14	15	16	17	18	19	20	File No.	Variables
2.90	3.44	1.25	1.78	0.63	1.23	2.31	2.27	2.76	1.44		Indexed-AAVA Cm ² /m ²
1.30	2.67	1.32	1.78	0.63	1.23	2.31	1.81	1.42	1.92		Indexed-EAVA Cm ² /m ²
0.44	0.77	1.00	1.00	1.00	1.00	1.00	0.79	0.51	1.00		EAVA/AAVA Ratio
0.31	2.28	2.82	2.11	1.77	2.62	3.50	2.56	2.42	2.79		Indexed-LVOTSA Cm ² /m ²
1.65	1.82	1.92	1.44	1.77	3.76	1.97	1.62	2.47	1.75		V _{max} - M/s
1.08	1.28	1.33	0.97	1.29	2.29	1.28	0.97	1.74	1.16		V _{mean} - M/s
11.02	12.89	14.57	8.42	12.69	56.56	15.68	10.64	24.46	12.34		P _{max} - mmHg
5.65	7.91	8.23	4.57	7.75	26.41	8.21	4.91	14.54	0.39		P _{mean} - mmHg
27.86	25.83	37.93	21.82	38.80	56.40	35.74	22.31	46.00	32.78		VTI-Cm
0.91	1.81	0.80	1.39	0.77	1.04	1.33	1.21	1.52	1.09		V _{max} - M/s
0.68	1.27	0.67	0.91	0.66	0.80	0.86	0.74	1.21	0.80		V _{mean} - M/s
3.52	12.08	2.61	7.72	2.36	4.33	7.07	5.94	9.36	4.86		P _{max} - mmHg
2.31	7.68	1.98	4.29	2.02	2.90	3.67	2.76	6.92	3.05		P _{mean} - mmHg
11.00	30.20	16.98	18.33	12.50	26.44	23.61	15.79	27.03	22.54		VTI-Cm
1.72	1.67	1.72	1.58	1.26	1.70	2.07	1.70	1.67	1.71		LVOT
1.61	1.79	1.31	1.51	1.10	1.20	1.87	1.60	1.78	1.23		AV-Ann
1.49	2.17	1.98	2.29	1.57	2.02	2.07	2.50	1.86	1.78		Vals-Sin
1.23	2.00	1.28	1.77	1.17	1.73	1.60	2.02	1.31	1.28		STJ
1.46	2.17	1.56	1.96	1.20	2.39	1.96	2.46	1.60	1.72		AAO
21	22	23	24	25	26	27	28	29	30	File no.	variable
2.54	3.86	2.36	2.14	0.83	0.53	1.65	2.53	1.94	3.17		Indexed-AAVA Cm ² /m ²
0.86	2.89	1.50	1.41	0.83	0.52	0.83	0.85	1.78	1.38		Indexed-EAVA Cm ² /m ²
0.33	0.88	0.63	0.66	1.00	0.99	0.44	0.33	0.93	0.43		EAVA/AAVA Ratio
2.66	3.14	1.66	1.68	2.21	1.60	1.54	2.61	1.76	2.50		Indexed-LVOTSA Cm ² /m ²
2.37	2.26	1.50	1.80	2.01	3.72	2.92	3.73	1.35	1.47		V _{max} - M/s
1.62	1.64	0.87	1.17	1.38	2.65	1.92	2.66	1.21	1.20		V _{mean} - M/s
22.59	20.54	9.02	13.04	16.18	55.41	34.22	55.36	14.89	8.71		P _{max} - mmHg
13.03	12.62	3.78	6.63	9.31	33.65	18.81	33.80	7.09	6.70		P _{mean} - mmHg
43.34	46.13	22.12	26.14	39.53	91.87	59.13	91.91	37.91	27.94		VTI-Cm
0.87	2.03	1.30	1.60	0.84	1.40	1.59	1.41	1.92	3.05		V _{max} - M/s
0.67	1.50	0.82	0.98	0.70	1.03	1.29	1.62	1.35	2.91		V _{mean} - M/s
3.07	16.54	6.84	10.24	2.82	7.90	10.26	7.90	14.67	45.98		P _{max} - mmHg
2.19	10.62	3.32	4.82	2.22	5.19	7.68	5.17	8.75	38.21		P _{mean} - mmHg
14.10	42.36	20.00	21.98	14.94	30.40	31.87	30.35	37.90	15.50		VTI-Cm
1.65	1.88	1.53	1.62	1.88	1.50	1.56	2.00	1.91	1.89		LVOT
1.61	1.91	1.82	1.83	1.69	1.73	1.73	1.97	1.99	2.13		AV-Ann
1.48	2.27	2.00	2.49	1.98	2.28	2.38	2.56	2.74	2.70		Vals-Sin
1.38	1.92	1.80	1.88	1.60	1.82	1.70	2.83	2.66	1.68		STJ
1.22	2.45	2.03	2.17	3.52	3.83	2.48	2.83	2.75	2.78		AAO

Table III. Maximum, Minimum & Mean of variables

Indexed AAVA cm^2/m^2	3.44	3.44	3.44
Indexed EAVA cm^2/m^2	3.15	3.15	3.15
EAVA/ AAVA Ratio	1.00	1.00	1.00
Indexed LVOTSA cm^2/m^2	3.63	3.63	3.63
AVPG _{max} mmHg	56.56	56.56	56.56
AVPG _{mean} mmHg	33.80	33.80	33.80
AV-VTI m/s	91.91	91.91	91.91
LVOT-VTI m/s	52.33	52.33	52.33
STJ-Dia Cm	2.83	2.83	2.83
AV-An-Dia Cm	2.13	2.13	2.13
Sin-Vals-Dia Cm	2.74	2.74	2.74
A-AO-Dia Cm	3.83	3.83	3.83
Variable			
Range	Maximum	Maximum	Maximum

AAO= Ascending aorta= Ascending aorta; AOVSA= AOV surface area = Aortic valve surface area;
 AAVA= Anatomic-AV area= Anatomic Aortic valve area;AOV= AV= Aortic valve; EAVA= Effective AV area = Effective aortic valve area; Ra= Ratio; LVOTSA= LVOT surface area = Left ventricular outflow tract surface area; LVOT = LV outflow tract; LV= Left ventricle; Cm^2 = Square centimeter; M^2 = Square meter; V max= Maximum velocity; V mean= Mean velocity; M/S= Meter per second;mmHg= Millimeter Hg (Mercury); VTI= Velocity time integral; P max= Maximum pressure gradient across AV; P mean= Mean pressure gradient across AV; AV-Ann= Aortic valve annulus; Sin-Vals= Valsalva sinus;STJ= Sinotubular junction; AAO= Ascending aorta; Cm= Centimeter; Dia= Diameter;

Table IV. Comparison of matched anatomic variables between patients and control cohort

AAO Diameter	STJ-SAR Diameter	Sinuses Diameter	AV-Annulus Diameter	No	Variables/ Cohort
3.15 cm/m^2	2.70 cm/m^2	3.30 cm/m^2	2.55 cm/m^2	30	Patients
2.86 cm/m^2	2.45 cm/m^2	3.15 cm/m^2	2.40 cm/m^2	30	Controls
10.13%	10.20%	4.75%	6.25%	30	Difference
P<0.05	P<0.01	NS	P<0.005	30	P value

Indexed-Matched= All diameters Indexed Per m^2 ; M^2 = Square meter; AV=Aortic Valve; Sinuses=Valsalva Sinuses; STJ-SAR=Sinotubular Junction- Supraaortic Ridge; AAO=Ascending Aorta; BAV=Bicuspid Aortic Valve

Discussion

The findings of our study support the hypothesis that BAV and aortic root dilation may share a common developmental defect. AS and AI are common in BAV. Similar to other obstructive defects of left heart, BAV is significantly more common in males. Even without AS or AI, murmurs and clicks are very common in BAV.

There are few studies about BAV in the literature. In 2003, Nistri et al. reported the results of their study and concluded that in BAV, aortic root dimensions are larger than control healthy subjects. These excess values of diameter for aortic valve annulus, sinuses of Valsalva, STJ and proximal AAO were 7.5%, 11.6%, 15% and 43.9%, respectively ³ (Table V).

Table V. Comparison of our study with others

Hemodynamic variables			Anatomic variables				Variables/ Studies
Normal	AI	AS	AAO	STJ	Sinuses	Annulus	
60%	36%	40%	10.13%	10.20%	4.75%	6.25%	Our study
23%	64%	13%	50-60%	57-79%	36-78%	9-59%	Rebecca et al ⁽²⁾
-	-	-	44%	15%	11.60%	7.50%	s. Nistri et al ⁽³⁾

Habn et al. reported that these excess of dimension for the above-mentioned 4 levels were 9-59%, 36-78%, 57-79% and 50-64%. The BAV was more common in males.² In another study by Paul, et al. they reported that BAV is the most common CHD and may be a genetic defect. BAV is a disease of the aortic root at all levels.¹

While our study supports the results of the other studies in other centers, it also shows differences in the dimensions and other values measured by us and others. Because other studies were done in adult subjects in contrast to ours, which was carried out in children and young patients, and because the aortic root dilatation, AS and AI are progressive, the percents of excess in aortic root dimension must be smaller in our study. According to our study and other studies we recommend:

- 1.Evaluation of all children with heart murmur and / or click to rule out BAV,
- 2.Follow-up of all patients with BAV for progression of AS, AI and other complications,
- 3.Screening of the patient's first-degree relatives for BAV by noninvasive screening and diagnostic tests such as transthoracic echocardiography, and

4. Evaluation of BAV cases for the co-existence of CHD.

Acknowledgments

We would like to thank the staff of pediatric echocardiography department of Shaheed Rajaie Heart Hospital for their collaboration.

References

- Paul WM, Fedak MD. Clinical and pathophysiological implications of a BAV. *Circulation* 106: 900; 2002.
- Hahn RT. Stenotic and functionally normal BAV. *JACC* 19 (2): 283-8, 1992.
- Nistri S. Aortic root dilation in young BAV. *Heart* 82: 1; 19-22, 2003.
- Allen HD. In: Moss and Adams' Heart Diseases in Infants, Children and Adolescents. 6th ed, Lippincott, Williams & Wilkins, Pennsylvania, USA, 2001.
- Snider AR. Echocardiography in Pediatric Heart Diseases. 2nd ed, Mosby Year Book, St. Louis, USA, 1997.
- Garson A, Jr. The Science and Practice of Pediatric Cardiology. 2nd ed, Williams and Wilkins, Pennsylvania, USA, 1998.
- Beppu S, Suzuki S, Matsuda. Rapidity of progression of aortic stenosis in patients with congenital BAV. *Am J Cardiol* 71: 322-327, 1993.
- Gersony WM, Hayes CJ, Driscoll DJ. Second natural history study of CHD Congenital Heart Defect *Circulation* 87: 152-165, 1993.
- Bonow RO, Carabello B, De Leon AC. ACC/AHA Guidelines for the management of patients with valvular heart disease. *J Am Coll Cardiol* 32: 1486-1588, 1998.
- McCrinkle BW. Independent predictors of immediate results of PTAV in children with Bicuspid Aortic Valve *Am J Cardiol* 77: 286-93, 1996.
- Kouchoukos NT, Davila-Roman VG, Spray TL. Replacement of Ao-Root in Bicuspid Aortic valve *N Engl J Med*: 330: 1-6, 1994.
- Phillips JR. Valvular hemodynamin in auto graft valve replacement (Ross procedural [Abstract]. *J Am Coll Cardiol* 33: 559, 1999.
- Roberts CS, Roberts WC. Dissection of the aorta in Bicuspid Aortic valve *J Am Coll Cardiol* 33(17): 712-716, 1991.
- Trivedi KR. Echocardiographic assessment of aortic root in children with BAV. *Heart* 78: 32, 1997.
- Hanh RT. Association of Aortic root dilation with BAV *J Am Coll Cardiol* 17: 1991.
- Hanh RT, Roman. Association of BAV with AI and aortic dilation (Case review). *J Am Coll Cardiology*, 19: 283-288, 1992.
- Levinson GE, Alpert JS. Aortic stenosis Valvular heart diseases. 3rd ed. Philadelphia, Lippincott Williams and Wilkins, 2000.
- Huntington K, Hunter AGW, Chan KL: A prospective study to assess the frequency of familial clustering of congenital BAV. *J Am Coll Cardiol* 30:1809-1812, 1997.
- Fedak PWM, Verma S, David TE: Clinical and pathophysiology implication of a BAV. *Circulation* 106: 900, 2002.
- Keane MG, Wiegers SE, Plappert T. BAVs are associated with aortic dilatation out of proportion to coexistence of valvular lesions. *Circulation* 102: III-35, 2000.
- Nataatmadja M. Abnormal extracellular matrix protein transport associated with increased apoptosis of vascular smooth muscle cells in Marfan syndrome and BAV, thoracic aortic aneurysm. *Circulation* 108:II-329, 2003.
- Ward C. Clinical significance of BAV. *Heart* 83: 81-5, 2000.
- Nistri S. Aortic root dilation in young men with normally functioning BAVs. *Heart*; 82: 19-22, 1999.
- Parai JL. Aortic medial changes associated with BAVs. *Can J Cardiol* 15: 1233-38, 1999.
- J.A, SOC echocardiography 16: 1201-3; 2003.